

A Class of Irreducible Modules for the Extended Affine Lie Algebra $\widetilde{\mathfrak{gl}}_l(\mathbb{C}_q)$

Ziting Zeng*

School of Mathematical Science, Beijing Normal University
Beijing, China 100875
zengzt@bnu.edu.cn

Abstract

We construct a class of modules for extended affine Lie algebra $\widetilde{\mathfrak{gl}}_l(\mathbb{C}_q)$ by using the free fields. A necessary and sufficient condition is given for those modules being irreducible.

1 Introduction

The representation theory of affine Kac-Moody Lie algebras has remarkable applications in many areas of mathematics and mathematical physics. Extended affine Lie algebras are a higher dimensional generalization of affine Kac-Moody Lie algebras first introduced by [H-KT] under the name of quasi-simple Lie algebras and systematically studied in [AABGP]. This family of newly developed Lie algebras includes toroidal Lie algebras and the central extensions of the matrix Lie algebras coordinated by quantum tori as examples. The study of representations for extended affine Lie algebras coordinated by quantum tori have drawn a lot of attentions recently via various module realizations. For instance, the representations via vertex operator construction was obtained in [BS], [G1], [G2], [BGT]. Fermionic and bosonic realizations were constructed in [G3] and [L]. Unitary representations were studied in [JK], [ER], [GZ1] and [Z]. A deformed version was given in [VV].

The Wakimoto's free fields construction provides a remarkable way to realize affine Kac-Moody Lie algebras (see [W1], [FF] and [W2]). In [GZ]

*The author gratefully acknowledges the grant support from NNSF of China No.10801010.

and [Z], this approach has been successfully used to construct the representation of $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$, the extended affine Lie algebra of type A coordinated by a quantum torus, for $l = 2, 3$. Motivated by [Z] and [GZ2], we shall construct in this paper a class of irreducible representations of the extended affine Lie algebra $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$ which contains $l = 2$ in [GZ] as its special case. We further give a necessary and sufficient condition for those modules being irreducible.

The paper is organized as follows. In Sect 2, we recall some notations and definitions. Our main theorem will appears in Sect 3 where we construct a class of modules for $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$ by using the free fields. Then we work out the condition of irreducibility for modules constructed in Sect 4.

2 Preliminary

Let q be a non-zero complex number. A quantum 2-torus associated to q is the unital associative $\mathbb{C}$ -algebra $\mathbb{C}_q[s^{\pm 1}, t^{\pm 1}]$ (or, simply $\mathbb{C}_q$) with generators $s^{\pm 1}, t^{\pm 1}$ and relations

$$ss^{-1} = s^{-1}s = tt^{-1} = t^{-1}t = 1 \text{ and } ts = qst.$$

Let d_s, d_t be the degree operators on $\mathbb{C}_q$ defined by

$$d_s(s^m t^n) = m s^m t^n, \quad d_t(s^m t^n) = n s^m t^n$$

for $m, n \in \mathbb{Z}$.

Let $\mathfrak{gl}_l$ be the general linear algebra, which is one dimensional central extension of the simple Lie algebra $\mathfrak{sl}_l$ of type A_{l-1} . Denote $\mathfrak{h}$ the Cartan subalgebra of $\mathfrak{gl}_l$ which is the set of the diagonal matrices, and $\dot{P} \subset \mathfrak{h}^*$ be the set of weight lattice. Its Borel subalgebra denoted as $\mathfrak{b}$ is the set of all upper triangular matrices.

We form a natural central extension of $\mathfrak{gl}_l \otimes \mathbb{C}_q$ as follows.

$$\widehat{\mathfrak{gl}_l(\mathbb{C}_q)} = \mathfrak{gl}_l \otimes \mathbb{C}_q \oplus \mathbb{C}c_s \oplus \mathbb{C}c_t$$

with Lie bracket

$$\begin{aligned} & [E_{ij} \otimes s^{m_1} t^{n_1}, E_{kn} \otimes s^{m_2} t^{n_2}] \\ &= \delta_{jk} q^{n_1 m_2} E_{in} \otimes s^{m_1+m_2} t^{n_1+n_2} - \delta_{in} q^{n_2 m_1} E_{kj} \otimes s^{m_1+m_2} t^{n_1+n_2} \\ &+ m_1 q^{n_1 m_2} \delta_{jk} \delta_{in} \delta_{m_1+m_2, 0} \delta_{n_1+n_2, 0} c_s + n_1 q^{n_1 m_2} \delta_{jk} \delta_{in} \delta_{m_1+m_2, 0} \delta_{n_1+n_2, 0} c_t \end{aligned}$$

for $m_1, m_2, n_1, n_2 \in \mathbb{Z}$, $1 \leq i, j, k, n \leq l$, where E_{ij} is the matrix whose (i, j) -entry is 1 and 0 elsewhere, and c_s and c_t are central elements of $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$.

The derivations d_s and d_t can be extended to derivations on $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$. Now we can define the semi-direct product of the Lie algebra $\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}$ and those derivations:

$$\widetilde{\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}} = \widehat{\mathfrak{gl}_l(\mathbb{C}_q)} \oplus \mathbb{C}d_s \oplus \mathbb{C}d_t.$$

The Lie algebra $\widetilde{\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}}$ is an extended affine Lie algebra of type A_{l-1} with nullity 2. (See [AABGP] and [BGK] for definitions).

So the Cartan subalgebra $\mathfrak{h}$ of $\widetilde{\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}}$ is

$$\mathfrak{h} = \dot{\mathfrak{h}} \oplus \mathbb{C}c_s \oplus \mathbb{C}c_t \oplus \mathbb{C}d_s \oplus \mathbb{C}d_t,$$

the Borel subalgebras $\mathfrak{b}$ is

$$\mathfrak{b} = \dot{\mathfrak{b}} \otimes \mathbb{C}_q \oplus \mathbb{C}c_s \oplus \mathbb{C}c_t \oplus \mathbb{C}d_s \oplus \mathbb{C}d_t$$

and denote $\mathfrak{n}_+ = \mathbb{C}\{E_{ij} \otimes \mathbb{C}_q, j > i \geq 1\}$ the nilpotent radical of $\mathfrak{b}$.

3 The module

In this section, we will construct the modules of $\widetilde{\widehat{\mathfrak{gl}_l(\mathbb{C}_q)}}$ by using the free fields.

Let

$$V = \mathbb{C}[x_i(m, n), i = 2, \dots, l, m, n \in \mathbb{Z}]$$

be a polynomial ring with infinitely many variables $x_i(m, n)$, we define the following operators:

$$\begin{aligned} e_{i1}(m_1, n_1) &= x_i(m_1, n_1), \quad i = 2, \dots, l; \\ e_{1i}(m_1, n_1) &= q^{-m_1 n_1} \mu \frac{\partial}{\partial x_i(-m_1, -n_1)} \\ &\quad - \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m', n') \in \mathbb{Z}^2}} q^{n_1 m' + n m_1 + n m'} x_2(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_2(m, n)} \frac{\partial}{\partial x_i(m', n')} \\ &\quad - \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m', n') \in \mathbb{Z}^2}} q^{n_1 m' + n m_1 + n m'} x_3(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_3(m, n)} \frac{\partial}{\partial x_i(m', n')} \\ &\quad - \dots \\ &\quad - \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m', n') \in \mathbb{Z}^2}} q^{n_1 m' + n m_1 + n m'} x_l(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_l(m, n)} \frac{\partial}{\partial x_i(m', n')}, \end{aligned}$$

$$\begin{aligned}
& \text{for } i = 2, \dots, l; \\
e_{ij}(m_1, n_1) &= \sum_{(m,n) \in \mathbb{Z}^2} q^{mn_1} x_i(m_1 + m, n_1 + n) \frac{\partial}{\partial x_j(m, n)}, \quad i, j = 2, \dots, l; \\
e_{11}(m_1, n_1) &= \mu \delta_{(m_1, n_1), (0,0)} - \sum_{(m,n) \in \mathbb{Z}^2} q^{nm_1} x_2(m_1 + m, n_1 + n) \frac{\partial}{\partial x_2(m, n)} \\
&\quad - \sum_{(m,n) \in \mathbb{Z}^2} q^{nm_1} x_3(m_1 + m, n_1 + n) \frac{\partial}{\partial x_3(m, n)} \\
&\quad - \dots \\
&\quad - \sum_{(m,n) \in \mathbb{Z}^2} q^{nm_1} x_l(m_1 + m, n_1 + n) \frac{\partial}{\partial x_l(m, n)} \\
D_1 &= \sum_{i=2}^l \sum_{(m,n) \in \mathbb{Z}^2} m x_i(m, n) \frac{\partial}{\partial x_i(m, n)} \\
D_2 &= \sum_{i=2}^l \sum_{(m,n) \in \mathbb{Z}^2} n x_i(m, n) \frac{\partial}{\partial x_i(m, n)}
\end{aligned}$$

Though some operators are infinite sums, they are well-defined when acting on V (as only finite summands left). Here we give our first theorem:

Theorem 3.1. *There is a Lie algebra homomorphism $\phi : \widetilde{\mathfrak{gl}_l(\mathbb{C}_q)} \rightarrow \mathfrak{gl}(V)$ given by*

$$\begin{aligned}
\phi(E_{ij} \otimes s^m t^n) &= e_{ij}(m, n), \quad i, j = 1, \dots, l, \\
\phi(d_s) &= D_1, \phi(d_t) = D_2, \phi(c_s) = \phi(c_t) = 0.
\end{aligned}$$

Hence V is a module of $\widetilde{\mathfrak{gl}_l(\mathbb{C}_q)}$.

Proof: We only need to check that ϕ preserves the brackets. The following formidable calculation is to check them case by case. For neatness we simplify $\sum_{\substack{(m,n) \in \mathbb{Z}^2 \\ (m',n') \in \mathbb{Z}^2}}$ or $\sum_{(m,n) \in \mathbb{Z}^2}$ as $\sum$.

If $i \neq j$

$$\begin{aligned}
& [e_{i1}(m_1, n_1), e_{1j}(m_2, n_2)] \\
&= [x_i(m_1, n_1), - \sum q^{n_2 m' + n m_2 + n m'} x_i(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_j(m', n')}] \\
&= \sum q^{n_2 m' + n_1 m_2 + n_1 m'} x_i(m_1 + m_2 + m', n_1 + n_2 + n') \frac{\partial}{\partial x_j(m', n')}
\end{aligned}$$

$$=q^{n_1 m_2} e_{ij}(m_1 + m_2, n_1 + n_2).$$

$$\begin{aligned}
& [e_{i1}(m_1, n_1), e_{1i}(m_2, n_2)] \\
&= [x_i(m_1, n_1), q^{-m_2 n_2} \mu \frac{\partial}{\partial x_i(-m_2, -n_2)}] \\
&+ [x_i(m_1, n_1), -\sum q^{n_2 m' + n m_2 + n m'} x_2(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_2(m, n)} \frac{\partial}{\partial x_i(m', n')}] \\
&+ \dots \\
&+ [x_i(m_1, n_1), -\sum q^{n_2 m' + n m_2 + n m'} x_i(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_i(m', n')}] \\
&+ \dots \\
&+ [x_i(m_1, n_1), -\sum q^{n_2 m' + n m_2 + n m'} x_l(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_l(m, n)} \frac{\partial}{\partial x_i(m', n')}] \\
&= -q^{-m_2 n_2} \mu \delta_{(m_1+m_2, n_1+n_2), (0,0)} + \sum q^{n_2 m_1 + n m_2 + n m_1} x_2(m_2 + m + m_1, n_2 + n + n_1) \frac{\partial}{\partial x_2(m, n)} \\
&+ \dots \\
&+ \sum q^{n_2 m_1 + n m_2 + n m_1} x_i(m_2 + m + m_1, n_2 + n + n_1) \frac{\partial}{\partial x_i(m, n)} \\
&+ \sum q^{n_2 m' + n_1 m_2 + n_1 m'} x_i(m_2 + m_1 + m', n_2 + n_1 + n') \frac{\partial}{\partial x_i(m', n')} \\
&+ \dots \\
&+ \sum q^{n_2 m_1 + n m_2 + n m_1} x_l(m_2 + m + m_1, n_2 + n + n_1) \frac{\partial}{\partial x_l(m, n)} \\
&= -q^{m_1 n_2} \mu \delta_{(m_1+m_2, n_1+n_2), (0,0)} + q^{n_2 m_1} (\sum q^{n(m_2+m_1)} x_2(m_2 + m + m_1, n_2 + n + n_1) \frac{\partial}{\partial x_2(m, n)} \\
&\dots \\
&+ \sum q^{n(m_2+m_1)} x_l(m_2 + m + m_1, n_2 + n + n_1) \frac{\partial}{\partial x_l(m, n)} \\
&+ q^{n_1 m_2} \sum q^{(n_2+n_1)m'} x_i(m_2 + m_1 + m', n_2 + n_1 + n') \frac{\partial}{\partial x_i(m', n')} \\
&= -q^{n_2 m_1} e_{11}(m_1 + m_2, n_1 + n_2) + q^{n_1 m_2} e_{ii}(m_1 + m_2, n_1 + n_2).
\end{aligned}$$

$$[e_{i1}(m_1, n_1), e_{jk}(m_2, n_2)]$$

$$\begin{aligned}
&= [x_i(m_1, n_1), \sum q^{mn_2} x_j(m_2 + m, n_2 + n) \frac{\partial}{\partial x_k(m, n)}] \\
&= -\delta_{ki} q^{m_1 n_2} x_j(m_1 + m_2, n_1 + n_2) \\
&= -\delta_{ki} q^{m_1 n_2} e_{j1}(m_1 + m_2, n_1 + n_2).
\end{aligned}$$

$$\begin{aligned}
&[e_{i1}(m_1, n_1), e_{11}(m_2, n_2)] \\
&= [x_i(m_1, n_1), -\sum q^{nm_2} x_i(m_2 + m, n_2 + n) \frac{\partial}{\partial x_i(m, n)}] \\
&= q^{n_1 m_2} x_i(m_1 + m_2, n_1 + n_2) \\
&= q^{n_1 m_2} e_{i1}(m_1 + m_2, n_1 + n_2).
\end{aligned}$$

$$\begin{aligned}
&[e_{1i}(m_1, n_1), e_{1i}(m_2, n_2)] \\
&= [q^{-m_1 n_1} \mu \frac{\partial}{\partial x_i(-m_1, -n_1)} \\
&\quad - \sum q^{n_1 m' + n m_1 + n m'} x_j(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')}, \\
&\quad q^{-m_2 n_2} \mu \frac{\partial}{\partial x_i(-m_2, -n_2)} \\
&\quad - \sum q^{n_2 m' + n m_2 + n m'} x_j(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')}] \\
&= -\mu q^{m_1 n_1} \sum_{\substack{-m_1 = m_2 + m + m' \\ -n_1 = n_2 + n + n'}} q^{n_2 m' + n m_2 + n m'} \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&\quad + \mu q^{-m_2 n_2} \sum_{\substack{-m_2 = m_1 + m + m' \\ -n_2 = n_1 + n + n'}} q^{n_1 m' + n m_1 + n m'} \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&\quad + \sum q^{n_1(m_2 + m'' + m') + n m_1 + n(m_2 + m'' + m') + n_2 m' + n'' m_2 + n'' m'} \\
&\quad \cdot x_j(m_1 + m + m_2 + m'' + m', n_1 + n + n_2 + n'' + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \frac{\partial}{\partial x_i(m', n')} \\
&\quad + \sum q^{n_1 m' + (n_2 + n + n'') m_1 + (n_2 + n + n'') m' + n_2 m'' + n m_2 + n m''} \\
&\quad \cdot x_j(m_1 + m + m_2 + m'' + m', n_1 + n + n_2 + n'' + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \frac{\partial}{\partial x_i(m', n')} \\
&\quad - \sum q^{n_2(m_1 + m'' + m') + n m_2 + n(m_1 + m'' + m') + n_1 m' + n'' m_1 + n'' m'} \\
&\quad \cdot x_j(m_1 + m + m_2 + m'' + m', n_1 + n + n_2 + n'' + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \frac{\partial}{\partial x_i(m', n')}
\end{aligned}$$

$$\begin{aligned}
& - \sum q^{n_2 m' + (n_1 + n + n'') m_2 + (n_1 + n + n'') m' + n_1 m'' + n m_1 + n m''} \\
& \cdot x_j(m_1 + m + m_2 + m'' + m', n_1 + n + n_2 + n'' + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \frac{\partial}{\partial x_i(m', n')} \\
& = 0.
\end{aligned}$$

If $i \neq j$,

$$\begin{aligned}
& [e_{1i}(m_1, n_1), e_{1j}(m_2, n_2)] \\
& = [- \sum q^{n_1 m' + n m_1 + n m'} x_k(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')}, \\
& \quad - \sum q^{n_2 m' + n m_2 + n m'} x_k(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_j(m', n')}] \\
& \quad + [q^{-m_1 n_1} \mu \frac{\partial}{\partial x_i(-m_1, -n_1)}, \\
& \quad - \sum q^{n_2 m' + n m_2 + n m'} x_i(m_2 + m + m', n_2 + n + n') \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_j(m', n')}] \\
& \quad + [- \sum q^{n_1 m' + n m_1 + n m'} x_j(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')}, \\
& \quad q^{-m_2 n_2} \mu \frac{\partial}{\partial x_j(-m_2, -n_2)}] \\
& = \sum q^{n_1(m_2 + m'' + m') + n m_1 + n(m_2 + m'' + m') + n_2 m' + n'' m_2 + n'' m'} \\
& \quad \cdot x_k(m_1 + m_2 + m + m' + m'', n_1 + n_2 + n + n' + n'') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \frac{\partial}{\partial x_j(m', n')} \\
& \quad + \sum q^{n_1 m' + (n_2 + n + n'') m_1 + (n_2 + n + n'') m' + n_2 m'' + n m_2 + n m''} \\
& \quad \cdot x_k(m_1 + m_2 + m + m' + m'', n_1 + n_2 + n + n' + n'') \frac{\partial}{\partial x_i(m', n')} \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_j(m'', n'')} \\
& \quad - \sum q^{n_2(m_1 + m' + m'') + n m_2 + n(m_1 + m' + m'') + n_1 m' + n'' m_1 + n'' m'} \\
& \quad \cdot x_k(m_1 + m_2 + m + m' + m'', n_1 + n_2 + n + n' + n'') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_j(m'', n'')} \frac{\partial}{\partial x_i(m', n')} \\
& \quad - \sum q^{n_2 m' + (n_1 + n + n'') m_2 + (n_1 + n + n'') m' + n_1 m'' + n m_1 + n m''} \\
& \quad \cdot x_k(m_1 + m_2 + m + m' + m'', n_1 + n_2 + n + n' + n'') \frac{\partial}{\partial x_j(m', n')} \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m'', n'')} \\
& \quad - \mu q^{-m_1 n_1} \sum_{\substack{-m_1 = m_2 + m + m' \\ -n_1 = n_2 + n + n'}} q^{n_2 m' + n m_2 + n m'} \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_j(m', n')}
\end{aligned}$$

$$\begin{aligned}
& + \mu q^{-m_2 n_2} \sum_{\substack{-m_2=m_1+m+m' \\ -n_2=n_1+n+n'}} q^{n_1 m' + n m_1 + n m'} \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& = 0
\end{aligned}$$

as the first and the fourth canceled each other and the second one and the third are negative to each other; and the last two kill each other according to the calculation of the case $[e_{1i}(m_1, n_1), e_{1i}(m_2, n_2)]$.

$$\begin{aligned}
& [e_{1i}(m_1, n_1), e_{ik}(m_2, n_2)] \\
& = q^{-m_1 n_1} \mu q^{(-m_1 - m_2) n_2} \frac{\partial}{\partial x_k(-m_1 - m_2, -n_1 - n_2)} \\
& \quad - \sum q^{n_1(m_2+m') + n m_1 + n(m_2+m') + m' n_2} \\
& \quad \cdot x_2(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_2(m, n)} \frac{\partial}{\partial x_k(m', n')} \\
& \quad - \dots \\
& \quad - \sum q^{n_1(m_2+m') + n m_1 + n(m_2+m') + m' n_2} \\
& \quad \cdot x_l(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_l(m, n)} \frac{\partial}{\partial x_k(m', n')} \\
& \quad - \sum q^{n_1 m' + (n_2 + n) m_1 + (n_2 + n) m' + m n_2} \\
& \quad \cdot x_i(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_i(m', n')} \frac{\partial}{\partial x_k(m, n)} \\
& \quad + \sum q^{(m_1 + m + m') n_2 + n_1 m' + n m_1 + n m'} \\
& \quad \cdot x_i(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& = q^{n_1 m_2} e_{1k}(m_1 + m_2, n_1 + n_2).
\end{aligned}$$

For $i \neq k$:

$$\begin{aligned}
& [e_{1i}(m_1, n_1), e_{kk}(m_2, n_2)] \\
& = [- \sum q^{n_1 m' + n m_1 + n m'} x_k(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')}, \\
& \quad \sum q^{m n_2} x_k(m_2 + m, n_2 + n) \frac{\partial}{\partial x_k(m, n)}] \\
& = - \sum q^{n_1 m' + (n_2 + n) m_1 + (n_2 + n) m' + m n_2}
\end{aligned}$$

$$\begin{aligned}
& \cdot x_k(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_i(m', n')} \frac{\partial}{\partial x_k(m, n)} \\
& + \sum q^{(m_1+m+m')n_2+n_1m'+nm_1+nm'} \\
& \cdot x_k(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& = 0.
\end{aligned}$$

For $i \neq k$ and $k \neq j$:

$$\begin{aligned}
& [e_{1i}(m_1, n_1), e_{kj}(m_2, n_2)] \\
& = [- \sum q^{n_1m'+nm_1+nm'} x_k(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& \quad - \sum q^{n_1m'+nm_1+nm'} x_j(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& \quad \sum q^{nm_2} x_k(m_2 + m, n_2 + n) \frac{\partial}{\partial x_j(m, n)}] \\
& = - \sum q^{n_1m'+(n_2+n)m_1+(n_2+n)m'+mn_2} \\
& \quad \cdot x_k(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_i(m', n')} \frac{\partial}{\partial x_j(m, n)} \\
& \quad + q^{(m_1+m+m')n_2+n_1m'+nm_1+nm'} \\
& \quad \cdot x_k(m_1 + m_2 + m + m', n_1 + n_2 + n + n') \frac{\partial}{\partial x_j(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& = 0.
\end{aligned}$$

$$\begin{aligned}
& [e_{1i}(m_1, n_1), e_{11}(m_2, n_2)] \\
& = [q^{-m_1n_1} \mu \frac{\partial}{\partial x_i(-m_1, -n_1)}, - \sum q^{nm_2} x_k(m_2 + m, n_2 + n) \frac{\partial}{\partial x_k(m, n)}] \\
& \quad + \sum_{k \neq i} [- \sum q^{n_1m'+nm_1+nm'} x_k(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& \quad - \sum q^{nm_2} x_k(m_2 + m, n_2 + n) \frac{\partial}{\partial x_k(m, n)}] \\
& \quad + \sum_{k=2}^n [- \sum q^{n_1m'+nm_1+nm'} x_k(m_1 + m + m', n_1 + n + n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
& \quad - \sum q^{nm_2} x_i(m_2 + m, n_2 + n) \frac{\partial}{\partial x_i(m, n)}]
\end{aligned}$$

$$\begin{aligned}
&= -q^{n_2 m_1} \mu q^{-(m_1+m_2)(n_1+n_2)} \frac{\partial}{\partial x_i(-(m_1+m_2), -(n_1+n_2))} \\
&\quad + \sum_{k \neq i} \left(\sum q^{n_1 m' + (n_2+n)m_1 + (n_2+n)m' + nm_2} \right. \\
&\quad \quad \cdot x_k(m_1+m_2+m+m', n_1+n_2+n+n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&\quad - \sum q^{(n_1+n+n')m_2 + n_1 m' + nm_1 + nm'} \\
&\quad \quad \cdot x_k(m_1+m_2+m+m', n_1+n_2+n+n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \Big) \\
&\quad + \sum q^{n_1 m' + (n_2+n)m_1 + (n_2+n)m' + nm_2} \\
&\quad \quad \cdot x_i(m_1+m_2+m+m', n_1+n_2+n+n') \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&\quad - \sum q^{(n_1+n+n')m_2 + n_1 m' + nm_1 + nm'} \\
&\quad \quad \cdot x_i(m_1+m_2+m+m', n_1+n_2+n+n') \frac{\partial}{\partial x_i(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&\quad + \sum_{k=2}^n \sum q^{n_1(m'+m_2) + nm_1 + n(m'+m_2) + n'm_2} \\
&\quad \quad \cdot x_k(m_1+m+m'+m_2, n_1+n+n'+n_2) \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&= -q^{n_2 m_1} \mu q^{-(m_1+m_2)(n_1+n_2)} \frac{\partial}{\partial x_i(-(m_1+m_2), -(n_1+n_2))} \\
&\quad + q^{n_2 m_1} \sum_{k \neq i} \sum q^{n_1 m' + nm_1 + (n_2+n)m' + nm_2} \\
&\quad \quad \cdot x_k(m_1+m_2+m+m', n_1+n_2+n+n') \frac{\partial}{\partial x_k(m, n)} \frac{\partial}{\partial x_i(m', n')} \\
&= -q^{n_2 m_1} e_{1i}(m_1+m_2, n_1+n_2).
\end{aligned}$$

If $1 < i, j, k, r \leq n$,

$$\begin{aligned}
&[e_{ij}(m_1, n_1), e_{kr}(m_2, n_2)] \\
&= [\sum q^{mn_1} x_i(m_1+m, n_1+n) \frac{\partial}{\partial x_j(m, n)}, \sum q^{mn_2} x_k(m_2+m, n_2+n) \frac{\partial}{\partial x_r(m, n)}] \\
&= \delta_{jk} \sum q^{(m_2+m)n_1 + mn_2} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_r(m, n)}
\end{aligned}$$

$$\begin{aligned}
& -\delta_{ir} \sum q^{(m_1+m)n_2+mn_1} x_k(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_j(m, n)} \\
& = \delta_{jk} q^{m_2 n_1} \sum q^{m(n_1+n_2)} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_r(m, n)} \\
& \quad - \delta_{ir} q^{m_1 n_2} \sum q^{m(n_1+n_2)} x_k(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_j(m, n)} \\
& = \delta_{jk} q^{m_2 n_1} e_{ir}(m_1+m_2, n_1+n_2) - \delta_{ir} q^{m_1 n_2} e_{kj}(m_1+m_2, n_1+n_2).
\end{aligned}$$

For $i \neq 1$,

$$\begin{aligned}
& [e_{ii}(m_1, n_1), e_{11}(m_2, n_2)] \\
& = [\sum q^{mn_1} x_i(m_1+m, n_1+n) \frac{\partial}{\partial x_i(m, n)}, -\sum q^{nm_2} x_i(m_2+m, n_2+n) \frac{\partial}{\partial x_i(m, n)}] \\
& = -\sum q^{(m_2+m)n_1+nm_2} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \\
& \quad + \sum q^{(n_1+n)m_2+mn_1} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \\
& = 0.
\end{aligned}$$

If $i \neq j$,

$$\begin{aligned}
& [e_{ij}(m_1, n_1), e_{11}(m_2, n_2)] \\
& = [\sum q^{mn_1} x_i(m_1+m, n_1+n) \frac{\partial}{\partial x_j(m, n)}, \\
& \quad -\sum q^{nm_2} x_i(m_2+m, n_2+n) \frac{\partial}{\partial x_i(m, n)} - \sum q^{nm_2} x_j(m_2+m, n_2+n) \frac{\partial}{\partial x_j(m, n)}] \\
& = -\sum q^{(m_2+m)n_1+nm_2} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_j(m, n)} \\
& \quad + \sum q^{(n_1+n)m_2+mn_1} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_j(m, n)} \\
& = 0.
\end{aligned}$$

$$\begin{aligned}
& [e_{11}(m_1, n_1), e_{11}(m_2, n_2)] \\
& = \sum_{i=2}^l [-\sum q^{m_1 n} x_i(m_1+m, n_1+n) \frac{\partial}{\partial x_i(m, n)}, \\
& \quad -\sum q^{m_2 n} x_i(m_2+m, n_2+n) \frac{\partial}{\partial x_i(m, n)}]
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=2}^l \left(\sum q^{m_1(n_2+n)+m_2n} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \right. \\
&\quad \left. - \sum q^{m_2(n_1+n)+m_1n} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \right) \\
&= -q^{m_1n_2} \mu_{\delta(m_1+m_2, n_1+n_2), (0,0)} \\
&\quad + \sum_{i=2}^l \sum q^{m_1(n_2+n)+m_2n} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \\
&\quad + q^{m_2n_1} \mu_{\delta(m_1+m_2, n_1+n_2), (0,0)} \\
&\quad - \sum_{i=2}^l \sum q^{m_2(n_1+n)+m_1n} x_i(m_1+m_2+m, n_1+n_2+n) \frac{\partial}{\partial x_i(m, n)} \\
&= -q^{m_1n_2} e_{11}(m_1+m_2, n_1+n_2) + q^{m_2n_1} e_{11}(m_1+m_2, n_1+n_2).
\end{aligned}$$

For the proof of brackets involving D_i , we can refer to [Z] for details. $\square$

It is obvious that V is a module of $\widetilde{\mathfrak{gl}_l(\mathbb{C}_q)}$ generated by 1, and $\mathfrak{n}_+.1 = 0$. Hence V is the highest weight module of $\widetilde{\mathfrak{gl}_l(\mathbb{C}_q)}$, and its highest weight vector is 1, with the weight α , such that

$$\alpha(E_{11}) = \mu, \alpha(E_{ii}) = 0, i \neq 1; \alpha(c_j) = \alpha(d_j) = 0, j = s, t$$

Remark 3.1. *The module given in [GZ1] is a special case of this module for $l = 2$. The one in [Z] is in a similar way of choosing different Borel subalgebra.*

4 Irreducibility

Since the centers c_s, c_t act on V trivially, we can view V as a module of $\mathfrak{gl}_l \otimes \mathbb{C}_q$. It is a weight module with the Cartan subalgebra $\mathfrak{h}$, the highest weight is $\alpha|_{\mathfrak{h}}$, we still denote it by α .

We need to figure out the weight spaces of V . Before doing that, we introduce some notations which will be used later.

Let

$$(M, N) = \begin{pmatrix} (m_{21}, n_{21}) & \cdots & (m_{2k_2}, n_{2k_2}) \\ (m_{31}, n_{31}) & \cdots & (m_{3k_3}, n_{3k_3}) \\ \cdots & \cdots & \cdots \\ (m_{l1}, n_{l1}) & \cdots & (m_{lk_l}, n_{lk_l}) \end{pmatrix}$$

with $k_i \in \mathbb{Z}_{\geq 0}$, $(m_{ij}, n_{ij}) \in \mathbb{Z}^2$, and $m_{i1} \leq m_{i2} \leq \cdots \leq m_{ik_i}$ for $i = 2, \dots, l$. (M, N) is a form with $l - 1$ lines, k_i is the length of $(i - 1)$ -line, length is 0 means this line is empty.

$(M, N) = (M', N')$ if two forms are the same; we call (M, N) and (M', N') the same type if the length of every line in (M, N) and (M', N') are the same. We denote the monomial $v_{(M, N)}$ as (M, N) defined above as

$$v_{(M, N)} = x_2(m_{21}, n_{21}) \cdots x_2(m_{2k_2}, n_{2k_2}) \cdots x_l(m_{l1}, n_{l1}) \cdots x_l(m_{lk_l}, n_{lk_l})$$

If $k_i = 0, i = 2, \dots, l$, this is 1.

Since

$$e_{11}(0, 0)(v_{(M, N)}) = \mu - k_2 - \cdots - k_l, e_{ii}(0, 0)(v) = k_i, i \neq 2,$$

the weight of $v_{(M, N)}$ is

$$\beta(E_{11}) = \mu - k_2 - \cdots - k_l, \beta(E_{ii}) = k_i, i \neq 1.$$

Hence the weight space is spanned by all those monomials $v_{(M, N)}$ with (M, N) the same type.

Theorem 4.1. *V is irreducible module of $\mathfrak{gl}_l \otimes \mathbb{C}_q$ if and only if $\mu \neq 0$.*

Proof: It is obvious that if $\mu = 0$, V is reducible, with the largest submodule is the polynomials without constant term.

On the other hand, since this module is a highest weight module, we only need to show that there is no other highest weight vector except 1 if $\mu \neq 0$.

Suppose

$$v = \sum_{(M, N)} a_{(M, N)} v_{(M, N)}$$

be a highest weight vector, here is a finite sum, and for all (M, N) are the same type, (i.e v is in a weight space). Hence $\mathfrak{n}_+$ acts as 0 on v .

At first we consider the case $j > i \geq 2$:

$$\begin{aligned} e_{ij}(m, n) \cdot v = \sum_{(M, N)} a_{(M, N)} \sum_{r=1}^{k_j} q^{m_{jr}n} x_i(m + m_{jr}, n + n_{jr}) \cdot x_2(m_{21}, n_{21}) \cdots \\ x_2(m_{2k_2}, n_{2k_2}) \cdots x_j(\widehat{m_{jr}, n_{jr}}) \cdots x_l(m_{l1}, n_{l1}) \cdots x_l(m_{lk_l}, n_{lk_l}) \end{aligned}$$

$$= \sum_{(M', N')} a'_{(M', N')} v_{(M', N')} = 0,$$

here $\widehat{}$ means that this term is omitted (the same for the following). Suppose (M, N) is not empty in some $(j-1)$ -th line, if we choose (m, n) big enough, then in (M', N') , the last one in the $(i-1)$ -th line is

$$(m_{jk_j} + m, n_{jk_j} + n),$$

so if $a'_{(M', N')} = ca_{(M, N)}$ for some (M, N) , with c is a nonzero complex number, so $a'_{(M', N')} = 0$, hence $a_{(M, N)} = 0$.

Therefore for the highest weight vector, type of (M, N) is empty in the lines other than the first one, so we simplify $(M, N) = ((m_1, n_1), \dots, (m_k, n_k))$ in the following.

Suppose that the highest weight vector is

$$v = \sum_{(M, N) \in K} a_{(M, N)} x_2(m_1, n_1) \cdots x_2(m_k, n_k)$$

where K is a set of some $(M, N) = ((m_1, n_1), \dots, (m_k, n_k))$ with the same types.

In the following, we consider the action of $e_{12}(a, b)$. It is easy to check that if $e_{12}(a, b)v_{(M, N)}$ and $e_{12}(a, b)v_{(M', N')}$ are linearly dependent for all big enough $a, b \in \mathbb{Z}$, then $v_{(M, N)}$ and $v_{(M', N')}$ must satisfy

$$\sum_{i=1}^k m_i = \sum_{i=1}^k m'_i, \quad \sum_{i=1}^k n_i = \sum_{i=1}^k n'_i$$

So we can assume that

$$\sum_{i=1}^k m_i = m, \quad \sum_{i=1}^k n_i = n, \quad (M, N) = ((m_1, n_1), \dots, (m_k, n_k)) \in K$$

for some $m, n \in \mathbb{Z}$.

If $k = 1$, then we have $v = ax_2(m, n)$, hence

$$0 = e_{12}(-m, -n)v = aq^{-mn}\mu.$$

So we get $a = 0$.

If $k = 2$, then

$$v = \sum_{i \in J} C_{m_i, n_i} x_2(m_i, n_i) x_2(m - m_i, n - n_i)$$

where J is an index set. Now let $I = \{(m_i, n_i) \mid i \in J\}$. By symmetries, we can assume $m_i \leq \frac{m}{2}$ for $i \in J$ and if $(\frac{m}{2}, n_j) \in I$ then $n_j \leq \frac{n}{2}$.

Now for $a, b \in \mathbb{Z}$, we have

$$0 = e_{12}(a, b)v = Ax_2(m + a, n + b)$$

where

$$\begin{aligned} A = & \mu q^{-ab}(C_{-a, -b} + C_{m+a, n+b}) \\ & + \sum_{i \in J} C_{m_i, n_i} \left(q^{b(m-m_i)+an_i+n_i(m-m_i)} + q^{bm_i+a(n-n_i)+m_i(n-n_i)} \right) \end{aligned}$$

First of all, for b big enough and all $a \in \mathbb{Z}$, we have

$$\begin{aligned} 0 = & \sum_{i \in J} \left(C_{m_i, n_i} q^{an_i+n_i(m-m_i)} q^{b(m-m_i)} + C_{m_i, n_i} q^{a(n-n_i)+m_i(n-n_i)} q^{bm_i} \right) \\ = & \sum_{i \in J'} A_{m_i} q^{b(m-m_i)} + \sum_{i \in J''} B_{m_i} q^{bm_i} \end{aligned}$$

where $J' \subseteq J$ such that

$$\{m_i \mid i \in J'\} = \{m_i \mid i \in J\} \text{ and } m_i \neq m_j \text{ if } i \neq j \in J';$$

$J'' \subseteq J'$ such that

$$\{m_i \mid i \in J''\} \dot{\cup} \left\{ \frac{m}{2} \right\} = \{m_i \mid i \in J'\} \text{ if } \frac{m}{2} \in \{m_i \mid i \in J'\}$$

and $J' = J''$ otherwise; and where

$$A_{m_i} = \sum_{(m_i, k) \in I} C_{m_i, n_i} \left(q^{ak+k(m-m_i)} + \delta_{\frac{m}{2}, m_i} q^{a(n-k)+m_i(n-k)} \right) \text{ for } i \in J'$$

and

$$B_{m_i} = \sum_{(m_i, k) \in I} C_{m_i, n_i} q^{a(n-k)+(n-k)m_i} \text{ for } i \in J''.$$

If q is not a root of unity, then we have $q^{(m-m_i)}$, $i \in J'$ and q^{m_i} , $i \in J''$ are $|J'| + |J''|$ distinct nonzero complex numbers. Because of the nonvanishing of Vandermonde determinant, we have that $A_{m_i} = 0$, $i \in J'$ and $B_{m_i} = 0$, $i \in J''$ hold for all $a \in \mathbb{Z}$. Now we have

$$0 = \sum_{i \in J} \left(C_{m_i, n_i} q^{an_i+n_i(m-m_i)} q^{b(m-m_i)} + C_{m_i, n_i} q^{a(n-n_i)+m_i(n-n_i)} q^{bm_i} \right)$$

for all $a, b \in \mathbb{Z}$.

If q is a primitive L -th root of unity, then for b big enough and all $a \in \mathbb{Z}$, we have

$$\begin{aligned} 0 &= \sum_{i \in J} \left(C_{m_i, n_i} q^{an_i + n_i(m-m_i)} q^{b(m-m_i)} + C_{m_i, n_i} q^{a(n-n_i) + m_i(n-n_i)} q^{bm_i} \right) \\ &= \sum_{j=0}^{L-1} D_j q^{bj} \end{aligned}$$

where

$$D_j = \sum_{\substack{i \in J' \\ j \equiv m-m_i \pmod{L}}} A_{m_i} + \sum_{\substack{i \in J'' \\ j \equiv m_i \pmod{L}}} B_{m_i}.$$

Because of $q^i \neq q^j$ if $0 \leq i \neq j \leq L-1$, so we have $D_j = 0$ for all $0 \leq j \leq L-1$ and $a \in \mathbb{Z}$. Hence

$$0 = \sum_{i \in J} \left(C_{m_i, n_i} q^{an_i + n_i(m-m_i)} q^{b(m-m_i)} + C_{m_i, n_i} q^{a(n-n_i) + m_i(n-n_i)} q^{bm_i} \right)$$

for all $a, b \in \mathbb{Z}$.

Now for any nonzero complex number q and all $a, b \in \mathbb{Z}$, we have

$$A = \mu q^{-ab} (C_{-a, -b} + C_{m+a, n+b}) = 0.$$

If $(\frac{m}{2}, \frac{n}{2}) \in I$, taking $a = -\frac{m}{2}$ and $b = -\frac{n}{2}$, we get $C_{\frac{m}{2}, \frac{n}{2}} = 0$.

If $(\frac{m}{2}, \frac{n}{2}) \neq (m_i, n_i) \in I$, then $(m-m_i, n-n_i) \notin I$.

Setting $a = -m_i$ and $b = -n_i$, we have

$$0 = \mu q^{-m_i n_i} (C_{m_i, n_i} + C_{m-m_i, n-n_i}) = \mu q^{-m_i n_i} C_{m_i, n_i},$$

thus $C_{m_i, n_i} = 0$. Now we know $C_{m_i, n_i} = 0$ for all $i \in J$, i.e., $v = 0$.

If $k > 2$ and $|K| = \alpha$, then

$$v = \sum_{\beta=1}^{\alpha} C_{\beta} x_2(m_{\beta 1}, n_{\beta 1}) x_2(m_{\beta 2}, n_{\beta 2}) \cdots x_2(m_{\beta k}, n_{\beta k})$$

where

$$C_{\beta} \in \mathbb{C}, m_{\beta, k-1} + m_{\beta k} \geq m_{\gamma, k-1} + m_{\gamma k} \text{ if } 1 \leq \beta < \gamma \leq \alpha$$

and for the same $m_{\beta, k-1} + m_{\beta k}$, giving the similar ordering on $n_{\beta, k-1} + n_{\beta k}$.

Now we rewrite v as following

$$v = \sum_{\beta=1}^{\alpha} C_{\beta} v_{\beta} x_2(m_{\beta, k-1}, n_{\beta, k-1}) x_2(m_{\beta k}, n_{\beta k}).$$

Assume that

$$v_1 = v_2 = \cdots = v_r \neq v_{r+1}$$

and set

$$m_{1,k-1} + m_{1,k} = \cdots = m_{r,k-1} + m_{r,k} = m'$$

and

$$n_{1,k-1} + n_{1,k} = \cdots = n_{r,k-1} + n_{r,k} = n'.$$

Then

$$\begin{aligned} v &= v_1 \sum_{\beta=1}^l C_\beta x_2(m_{\beta,k-1}, n_{\beta,k-1}) x_2(m' - m_{\beta,k-1}, n' - n_{\beta,k-1}) + \text{other terms} \\ &= v_1 v' + \text{other terms} . \end{aligned}$$

Now we can easily see that $e_{12}(a, b)v' = 0$ for big enough b and all $a \in \mathbb{Z}$. By above discussions, we have $v' = 0$. Continuing this process, we will have $v = 0$.

Now, we must have $k = 0$. i.e., 1 is the only highest weight vector up to a scalar multiple. $\square$

Together with the actions of d_s and d_t on V , we can easily obtain that

Corollary 4.1. *V is an irreducible module of $\widetilde{\mathfrak{gl}_l(\mathbb{C}_q)}$ if and only if $\mu \neq 0$.*

If $\mu = 0$, it is obvious that the module is reducible, with the maximal submodule W which consists of the polynomials without constant term. Here we can get a parallel result of Thm 2.4 [GZ2].

Acknowledgment

I am grateful to Professor Yun Gao for his encouragement and stimulating discussions. I would also like to thank Dr. Hongjia Chen for some helpful comments on this work.

References

- [AABGP] B. N. Allison, S. Azam, S. Berman, Y. Gao, A. Pianzola, *Extended affine Lie algebras and their root systems*, Memoir. Amer. Math. Soc. 126 (1997) no. 605
- [BGK] S. Berman, Y. Gao, Y. Krylyuk, *Quantum tori and the structure of elliptic quasi-simple Lie algebras*, J. Funct. Anal. 135 (1996) 339–389

- [BGT] S. Berman, Y. Gao, S. Tan, *A unified view of some vertex operator constructions*, Israel Journal of Mathematics 134 (2003) 29–60
- [BS] S. Berman and J. Szmigielski, *Principal realization for extended affine Lie algebra of type sl_2 with coordinates in a simple quantum torus with two variables*, Contemp. Math. 248 (1999) 39–67
- [ER] Eswara Rao, *Unitary modules for EALAs co-ordinatized by a quantum torus*, Comm. Algebra 31 (2003) 2245–2256
- [FF] B. Feigin, E. Frenkel, *Affine Kac-Moody algebras and semi-infinite flag manifolds*, Comm. Math. Phys. 128 (1990) 161–189
- [G1] Y. Gao, *Representations of extended affine Lie algebras coordinatized by certain quantum tori*, Compositio Mathematica 123 (2000) 1–25
- [G2] Y. Gao, *Vertex operators arising from the homogeneous realization for $\widehat{gl}_N$* , Comm. Math. Phys. 211 (2000) 745–777
- [G3] Y. Gao, *Fermionic and bosonic representations of the extended affine Lie algebra $\widehat{\mathfrak{gl}}_N(\mathbb{C}_q)$* , Canada Math Bull. 45 (2002) 623–633
- [GZ1] Y. Gao and Z. Zeng, *Hermitian representations of the extended affine Lie algebra $\widehat{\mathfrak{gl}}_2(\mathbb{C}_q)$* , Adv. Math. 207 (2006) 244–265
- [GZ2] Y. Gao and Z. Zeng, *Irreducible Wakimoto-like modules for the affine Lie algebra $\widehat{\mathfrak{gl}}_n$* , to appear
- [H-KT] R. Høegh-Krohn and B. Torresani, *Classification and construction of quasi-simple Lie algebras*, J. Funct. Anal. 89 (1990) 106–136
- [JK] H. P. Jakobsen and V. G. Kac, *A new class of unitarizable highest weight representations of infinite-dimensional Lie algebras. II*, J. Funct. Anal. 82 (1989) 69–90
- [L] M. Lau, *Bosonic and fermionic representations of Lie algebra central extensions*, Adv. Math. **194** (2005) 225–245
- [VV] M. Varagnolo and E. Vasserot, *Double-loop algebras and the Fock space*, Invent. Math. 133 (1998) 133–159
- [W1] M. Wakimoto, *Fock representations of the affine Lie algebra $A_1^{(1)}$* , Comm. Math. Phys. 104 (1986) 605–609

- [W2] M. Wakimoto, *Extended affine Lie algebras and a certain series of Hermitian representations*, Preprint (1985)
- [Z] Z. Zeng, *Unitary representations of the extended affine Lie algebra $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$* , Pacific J. Math. 233 (2007) 481–509